



NÁRODNÁ BANKA SLOVENSKA
EUROSYSTEM

NORMALITY TESTS FOR DEPENDENT DATA

ZACHARIAS PSARADAKIS,
MARIÁN VÁVRA

WORKING
PAPER

12/2017



© National Bank of Slovakia

www.nbs.sk

Imricha Karvaša 1

813 25 Bratislava

research@nbs.sk

November 2017

ISSN 1337-5830

The views and results presented in this paper are those of the authors and do not necessarily represent the official opinions of the National Bank of Slovakia.

All rights reserved.



Normality Tests for Dependent Data¹

Working paper NBS

Zacharias Psaradakis, Marián Vávra²

Abstract

The paper considers the problem of testing for normality of the one-dimensional marginal distribution of a strictly stationary and weakly dependent stochastic process. The possibility of using an autoregressive sieve bootstrap procedure to obtain critical values and P -values for normality tests is explored. The small-sample properties of a variety of tests are investigated in an extensive set of Monte Carlo experiments. The bootstrap version of the classical skewness–kurtosis test is shown to have the best overall performance in small samples.

JEL classification: C12, C15, C32

Key words: Autoregressive sieve bootstrap, Normality test, Weak dependence

Downloadable at <http://www.nbs.sk/en/publications-issued-by-the-nbs/working-papers>

¹The authors are grateful to Antonio Trujillo-Ortiz for making some of his Matlab code available.

²Zacharias Psaradakis, Department of Economics, Mathematics and Statistics, Birkbeck, University of London.
Marián Vávra, Research Department of the NBS.



1. INTRODUCTION

The problem of testing whether a sample of observations comes from a Gaussian distribution has attracted considerable attention over the years. This is not perhaps surprising in view of the fact that normality is a common maintained assumption in a wide variety of statistical procedures, including estimation, inference, and forecasting procedures. In the context of model building, a test for normality is often a useful diagnostic for assessing whether a particular type of stochastic model may provide an appropriate characterization of the data (for instance, non-linear models are unlikely to be an adequate approximation to a time series having a Gaussian one-dimensional marginal distribution). Normality tests may also be useful in evaluating the validity of different hypotheses and models to the extent that the latter rely on or imply Gaussianity, as is the case, for example, with some option pricing, asset pricing, and dynamic stochastic general equilibrium models found in the economics and finance literature. Other examples where normality or otherwise of the marginal distribution is of interest, include value-at-risk calculations (e.g., Cotter (2007)), and copula-based modelling for multivariate time series with the marginal distribution and the copula function being specified separately. Kilian and Demiroglu (2000) and Bontemps and Meddahi (2005) give further examples where testing for normality is of interest.

Although most of the literature on tests for normality has focused on the case of independent and identically distributed (i.i.d.) observations (see Thode (2002) for an extensive review), a number of tests which are valid for dependent data have also been proposed. These include tests based on empirical standardized cumulants (Lobato and Velasco (2004), Bai and Ng (2005)), moment conditions of various types (e.g., Epps (1987), Moulines and Choukri (1996), Bontemps and Meddahi (2005)), the bispectral density function (e.g., Hinich (1982), Nusrat and Harvill (2008), Berg et al. (2010)), and the empirical distribution function (Psaradakis and Vávra (2017)). Unlike normality tests for i.i.d. observations, whose finite-sample behaviour has been extensively studied (see, inter alia, Baringhaus et al. (1989), Romão et al. (2010), and Yap and Sim (2011)), a similar comparison, across a common set of data-generating mechanisms, of tests designed for dependent data is not currently available in the literature.

Our aim in this paper is twofold. First, we wish to investigate the small-sample size and power properties of tests for normality of the one-dimensional marginal distribution of a strictly stationary time series. The tests under consideration are some of those mentioned in the previous paragraph, as well as tests that rely on the empirical characteristic function of the data. Second, since in the presence of serial dependence conventional large-sample approximations to the null distributions of some of the test statistics under consideration are inaccurate, unknown, or depend on the correlation structure of the data in complicated ways, we wish to investigate the possibility of using bootstrap resampling to implement tests of normality. More specifically, we consider estimating the null sampling distributions of the test statistics of interest by means

of the so-called autoregressive sieve bootstrap, and thus obtain P -values and/or critical values for normality tests. The bootstrap method is based on the idea of approximating the data-generating mechanism by an autoregressive sieve, that is, a sequence of autoregressive models the order of which increases with the sample size (e.g., Kreiss (1992), Bühlmann (1997)). Bootstrap-based normality tests are straightforward to implement and, as our simulation experiments demonstrate, offer significant improvements over asymptotic tests, that is, tests that use critical values from the large-sample null distributions of the relevant test statistics.

The remainder of the paper is organized as follows. Section 2 provides an overview of the normality tests of interest. Section 3 discusses how the autoregressive sieve bootstrap may be used to implement tests for normality of dependent data. Section 4 examines the small-sample properties of asymptotic and bootstrap-based normality tests by means of Monte Carlo simulations. Section 5 summarizes and concludes.

2. PROBLEM AND TESTS

2.1 STATEMENT OF THE PROBLEM

Suppose that (X_1, X_2, \dots, X_n) are n consecutive observations from a strictly stationary, real-valued, discrete-time stochastic process $\mathcal{X} = \{X_t\}_{t=-\infty}^{\infty}$ having mean $\mu_X = \mathbb{E}(X_t)$ and variance $\sigma_X^2 = \mathbb{E}[(X_t - \mu_X)^2] > 0$. It is assumed that \mathcal{X} is weakly dependent, in the sense that its autocovariance sequence decays towards zero sufficiently fast so that the series $\sum_{\tau=0}^{\infty} \text{Cov}(X_t, X_{t-\tau})$ converges absolutely (and, consequently, \mathcal{X} has a continuous and bounded spectral density). The problem of interest is to test the composite null hypothesis that the one-dimensional marginal distribution of \mathcal{X} is Gaussian, that is,

$$\mathcal{H}_0 : (X_t - \mu_X)/\sigma_X \sim \mathcal{N}(0, 1), \quad (1)$$

where a tilde ' \sim ' means 'is distributed as'. The alternative hypothesis is that the distribution of X_t is non-Gaussian.

2.2 TESTS BASED ON SKEWNESS AND KURTOSIS

Bowman and Shenton (1975) and Jarque and Bera (1987) proposed a test for normality based on the empirical standardized third and fourth cumulants, exploiting the fact that for a normal distribution all cumulants of order higher than the second are zero. The test statistic is given by

$$JB = \frac{n\hat{\mu}_3^2}{6\hat{\mu}_2^3} + \frac{n(\hat{\mu}_4 - 3\hat{\mu}_2^2)^2}{24\hat{\mu}_2^4}, \quad (2)$$

where, for an integer $r \geq 2$, $\hat{\mu}_r = (1/n) \sum_{t=1}^n (X_t - \bar{X})^r$ and $\bar{X} = (1/n) \sum_{t=1}^n X_t$. For Gaussian i.i.d. data, JB is approximately χ_2^2 distributed for large n . Although a test which rejects when JB exceeds an appropriate quantile of the χ_2^2 distribution is clearly not guaranteed to have correct asymptotic level in the presence of serial dependence, it is arguably the most popular normality test in the literature and is available in many statistical and econometric packages (e.g., EViews, Matlab, Stata). It will, thus, serve as a benchmark for comparisons in our study.

Bai and Ng (2005) developed a related test which allows for weak dependence in the data. The test is based on the statistic

$$BN = \frac{n\hat{\mu}_3^2}{\hat{\zeta}_3\hat{\mu}_2^3} + \frac{n(\hat{\mu}_4 - 3\hat{\mu}_2^2)^2}{\hat{\zeta}_4\hat{\mu}_2^4}, \quad (3)$$

where $\hat{\zeta}_3$ and $\hat{\zeta}_4$ are consistent estimators of the asymptotic variance of $\sqrt{n}\hat{\mu}_2^{-3/2}\hat{\mu}_3$ and $\sqrt{n}\hat{\mu}_2^{-2}(\hat{\mu}_4 - 3\hat{\mu}_2^2)$, respectively. Following Bai and Ng (2005), $\hat{\zeta}_3$ and $\hat{\zeta}_4$ are constructed using a non-parametric kernel estimator of the relevant long-run covariance matrices; the triangular Bartlett kernel and a data-dependent bandwidth, selected according to the method of Andrews (1991), are used.

An alternative test, also based on skewness and kurtosis, was proposed by Lobato and Velasco (2004). The test statistic is defined as

$$LV = \frac{n\hat{\mu}_3^2}{6\hat{G}_3} + \frac{n(\hat{\mu}_4 - 3\hat{\mu}_2^2)^2}{24\hat{G}_4}, \quad (4)$$

where $\hat{G}_r = \sum_{\tau=1-n}^{n-1} \hat{\gamma}_\tau^r$ for $r = 3, 4$ and $\hat{\gamma}_\tau = (1/n) \sum_{t=|\tau|+1}^n (X_t - \bar{X})(X_{t-|\tau|} - \bar{X})$ for $\tau = 0, \pm 1, \dots, \pm(n-1)$. An advantage of the test based on LV is that the estimators of the asymptotic variance of $\sqrt{n}\hat{\mu}_3$ and $\sqrt{n}(\hat{\mu}_4 - 3\hat{\mu}_2^2)$ used do not involve any kernel smoothing or truncation (in contrast to the estimators $\hat{\zeta}_3$ and $\hat{\zeta}_4$ used in the case of BN). If \mathcal{X} is a Gaussian process, BN and LV are approximately χ_2^2 distributed for large n .

2.3 TEST BASED ON MOMENT CONDITIONS

Bontemps and Meddahi (2005) proposed a test based on moment conditions implied by the characterization of the normal distribution given in Stein (1972). The test is based on the statistic

$$BM = \left(\frac{1}{\sqrt{n}} \sum_{t=1}^n \hat{\mathbf{g}}_t \right) \hat{\Sigma}^{-1} \left(\frac{1}{\sqrt{n}} \sum_{t=1}^n \hat{\mathbf{g}}_t' \right), \quad (5)$$

where $\hat{\mathbf{g}}_t = (h_3(Z_t), \dots, h_\ell(Z_t))$ for some integer $\ell \geq 3$, $Z_t = \{n\hat{\mu}_2/(n-1)\}^{-1/2}(X_t - \bar{X})$, and $\hat{\Sigma}$ is a consistent estimator of the long-run covariance matrix of $\{\hat{\mathbf{g}}_t\}$. Here, $h_m(\cdot)$ stands for the normalized Hermite polynomial of degree m , given by

$$h_m(x) = \sqrt{m!} \sum_{i=0}^{\lfloor m/2 \rfloor} \frac{(-1)^i x^{m-2i}}{i!(m-2i)!2^i}, \quad -\infty < x < \infty, \quad m = 0, 1, 2, \dots,$$

where $\lfloor a \rfloor$ denotes the largest integer not greater than a . Under \mathcal{H}_0 , BM is approximately $\chi_{\ell-2}^2$ distributed for large n .

As in the case of the BN statistic, $\hat{\Sigma}$ is constructed using a Bartlett-kernel estimator with a data-dependent bandwidth chosen by the method of Andrews (1991). In light of the relatively poor small-sample size properties of the test reported in Bontemps and Meddahi (2005) for dependent data when Hermite polynomials of degree higher than 4 are used, we set $\ell = 4$ in our implementation of the test.

2.4 TESTS BASED ON THE EMPIRICAL DISTRIBUTION FUNCTION

Psaradakis and Vávra (2017) considered a test based on the Anderson–Darling distance statistic involving the weighted quadratic distance of the empirical distribution function of the data from a Gaussian distribution function. Putting $Y_t = \hat{\mu}_2^{-1/2}(X_t - \bar{X})$, the test rejects \mathcal{H}_0 for large values of the statistic

$$\begin{aligned} AD &= n \int_{-\infty}^{\infty} \frac{\{\hat{F}_Y(y) - \Phi(y)\}^2}{\Phi(y)\{1 - \Phi(y)\}} d\Phi(y) \\ &= -n - \frac{1}{n} \sum_{t=1}^n (2t - 1) [\log \Phi(Y_{(t)}) + \log\{1 - \Phi(Y_{(n+1-t)})\}], \end{aligned} \quad (6)$$

where \hat{F}_Y is the empirical distribution function of (Y_1, \dots, Y_n) , $Y_{(1)} \leq \dots \leq Y_{(n)}$ are the order statistics of (Y_1, \dots, Y_n) , and Φ is the standard normal distribution function. In the sequel, we also consider tests which reject \mathcal{H}_0 for large values of the Cramér–von Mises statistic

$$CM = n \int_{-\infty}^{\infty} \{\hat{F}_Y(y) - \Phi(y)\}^2 d\Phi(y) = \frac{1}{12n} + \sum_{t=1}^n \left(\Phi(Y_{(t)}) - \frac{2t-1}{2n} \right)^2, \quad (7)$$

or the Kolmogorov–Smirnov statistic

$$\begin{aligned} KS &= \sqrt{n} \sup_{-\infty < y < \infty} |\hat{F}_Y(y) - \Phi(y)| \\ &= \sqrt{n} \max_{1 \leq t \leq n} \left\{ \frac{t}{n} - \Phi(Y_{(t)}), \Phi(Y_{(t)}) - \frac{t-1}{n}, 0 \right\}. \end{aligned} \quad (8)$$

Since the asymptotic null distributions of these statistics have a rather complicated structure in the case of a composite null hypothesis even under i.i.d. conditions (cf. Durbin (1973), Stephens (1976)), critical values and/or P -values for the tests will be obtained by a suitable bootstrap procedure. Stute et al. (1993), Babu and Rao (2004), and Kojadinovic and Yan (2012) also considered bootstrap-based approaches to testing composite hypotheses for i.i.d. data, while Psaradakis and Vávra (2017) examined the case of linear processes that may exhibit strong, weak, or negative dependence.

2.5 TEST BASED ON THE EMPIRICAL CHARACTERISTIC FUNCTION

Epps and Pulley (1983) proposed a class of tests based on the weighted quadratic distance of the empirical characteristic function of the data from its pointwise limit under the null hypothesis of normality. Using the density of the $\mathcal{N}(0, 1/\hat{\mu}_2)$ distribution as a weight function (cf. Epps and Pulley (1983)), the test rejects for large values of the statistic

$$\begin{aligned} EP &= n \int_{-\infty}^{\infty} |\hat{\varphi}_Y(u) - \varphi(u)|^2 d\Phi(\hat{\mu}_2^{1/2}u) \\ &= \frac{n}{\sqrt{3}} + \frac{1}{n} \sum_{t=1}^n \sum_{s=1}^n \exp\left\{-\frac{1}{2}(Y_t - Y_s)^2\right\} - \sqrt{2} \sum_{t=1}^n \exp\left(-\frac{1}{4}Y_t^2\right) \end{aligned} \quad (9)$$

where $\hat{\varphi}_Y$ is the empirical characteristic function of (Y_1, \dots, Y_n) and φ is the characteristic function of Φ .

For Gaussian i.i.d. data, EP is asymptotically distributed as a weighted sum of infinitely many independent χ_1^2 random variables (Baringhaus and Henze (1988)). To the best of our knowledge, the asymptotic distribution of EP has not been established in the case of dependent data. We will use a bootstrap procedure to obtain critical and/or P -values for the test based on EP . We note that, in an i.i.d. context, Jiménez-Gamero et al. (2003) and Leucht and Neumann (2009) examined bootstrap-based inference for statistics (such as EP , AD , and CM) which may be expressed in the form of, or be approximated by, degenerate V -statistics involving estimated parameters. Leucht (2012) and Leucht and Neumann (2013) give related results for weakly dependent data.

2.6 TEST BASED ON THE BISPECTRUM

Hinich (1982) proposed a test for Gaussianity of a stochastic process based on its normalized bispectrum, exploiting the fact that the latter should be identically zero at all frequency pairs if the process is Gaussian. For some integer $k \geq 1$, the test used in the sequel is based on the statistic

$$H = \frac{2\pi n}{\delta M^2} \sum_{i=1}^k \frac{|\hat{f}_b(\omega_{1,i}, \omega_{2,i})|^2}{\hat{f}_s(\omega_{1,i})\hat{f}_s(\omega_{2,i})\hat{f}_s(\omega_{1,i} + \omega_{2,i})}, \quad (10)$$

where \hat{f}_s and \hat{f}_b are kernel-smoothed estimators of the spectral and bispectral density, respectively, of \mathcal{X} , M is a bandwidth parameter associated with \hat{f}_b , δ is a normalizing constant associated with \hat{f}_b , and $\Omega_k = \{(\omega_{1,i}, \omega_{2,i}), i = 1, \dots, k\}$ is a set of frequency pairs contained in $\Omega = \{(\omega_1, \omega_2) : 0 \leq \omega_1 \leq \pi, 0 \leq \omega_2 \leq \min\{\omega_1, 2(\pi - \omega_1)\}\}$ (see Berg et al. (2010) for more details). If \mathcal{X} is a Gaussian process, H is approximately χ_{2k}^2 distributed for large n .

In the sequel, we follow Berg et al. (2010) in taking Ω_k to be a subset of the grid of points contained in their Fig. 2, as well as in using a trapezoidal flat-top kernel function and a right-

pyramidal frustrum-shaped kernel function to construct the estimators \hat{f}_s and \hat{f}_b , respectively. A common bandwidth $M = \lfloor n^{1/3} \rfloor$ is used for \hat{f}_s and \hat{f}_b , and we set $k = \lfloor n/10 \rfloor$. We note that Berg et al. (2010) considered using an autoregressive sieve bootstrap approximation to the null distribution of H as an alternative to the χ_{2k}^2 large-sample approximation. Also note that, unlike the testing procedures discussed previously, which assess normality of the one-dimensional marginal distribution of \mathcal{X} , the test based on H assesses Gaussianity of the process \mathcal{X} (i.e., normality of all finite-dimensional distributions of \mathcal{X}).

3. BOOTSTRAP TESTS

Some of the normality tests described in Section 2, although asymptotically valid for dependent data, tend to suffer from substantial level distortion in finite samples (e.g., the bispectrum-based test). For some other tests, large-sample approximations to the null distribution of the relevant test statistic may not be straightforward to obtain because of the dependence in the data and the composite null hypothesis (e.g., tests based on the empirical distribution function or the empirical characteristic function). A convenient way of overcoming these difficulties is to use a suitable bootstrap procedure to approximate the sampling distribution of the test statistic of interest under the null hypothesis. In this paper, we propose to use the autoregressive sieve bootstrap to obtain such an approximation and construct bootstrap tests for normality.

The typical assumption underlying the autoregressive sieve bootstrap is that \mathcal{X} admits the representation

$$X_t - \mu_X = \sum_{j=1}^{\infty} \phi_j (X_{t-j} - \mu_X) + \varepsilon_t, \quad (11)$$

where $\{\phi_j\}_{j=1}^{\infty}$ is an absolutely summable sequence of real numbers and $\{\varepsilon_t\}_{t=-\infty}^{\infty}$ are i.i.d., real-valued, zero-mean random variables with finite, positive variance. The idea is to approximate (11) by a finite-order autoregressive model, the order of which increases simultaneously with the sample size at an appropriate rate, and use this model as the basis of a semi-parametric bootstrap scheme (see, inter alia, Kreiss (1992), Paparoditis (1996), Bühlmann (1997), Choi and Hall (2000), and Kreiss et al. (2011)).

Note that, under the additional assumption that the function $\phi(z) = 1 - \sum_{j=1}^{\infty} \phi_j z^j$ has no zeros inside or on the complex unit circle, (11) is equivalent to assuming that \mathcal{X} satisfies

$$X_t = \mu_X + \sum_{j=0}^{\infty} \psi_j \varepsilon_{t-j}, \quad \psi_0 = 1, \quad (12)$$

for some absolutely summable sequence of real numbers $\{\psi_j\}_{j=1}^{\infty}$. Hence, it is easy to see that the normality hypothesis (1) holds if ε_t is normally distributed. Conversely, (1) implies normality of the distribution of ε_t , which in turn implies Gaussianity of the causal linear process \mathcal{X} defined by (12).

Letting $S = S(X_1, \dots, X_n)$ be a statistic for testing the normality hypothesis (1), the algorithm used to obtain an autoregressive sieve bootstrap approximation to the null distribution of S can be described by the following steps:

- S1. For some integer $p \geq 1$ (chosen as a function of n so that p increases with n but at a slower rate), compute the p th order least-squares estimate $(\hat{\phi}_{p1}, \dots, \hat{\phi}_{pp})$ of the autoregressive coefficients for \mathcal{X} by minimizing

$$(n - 2p)^{-1} \sum_{t=p+1}^n \left\{ (X_t - \bar{X}) - \sum_{j=1}^p \phi_{pj}(X_{t-j} - \bar{X}) \right\}^2. \quad (13)$$

- S2. Given some initial values $(X_{-p+1}^*, \dots, X_0^*)$, generate bootstrap pseudo-observations (X_1^*, \dots, X_n^*) via the recursion

$$X_t^* - \bar{X} = \sum_{j=1}^p \hat{\phi}_{pj}(X_{t-j}^* - \bar{X}) + \hat{\sigma}_p \varepsilon_t^*, \quad t = 1, 2, \dots, \quad (14)$$

where $\hat{\sigma}_p^2$ is the minimum value of (13) and $\{\varepsilon_t^*\}$ are independent random variables each having the $\mathcal{N}(0, 1)$ distribution. Define the bootstrap analogue of S by the plug-in rule as $S^* = S(X_1^*, \dots, X_n^*)$ (i.e., by applying the definition of S to the bootstrap pseudo-data).

- S3. Repeat step S2 independently B times to obtain a collection of B replicates (S_1^*, \dots, S_B^*) of S^* . The empirical distribution of (S_1^*, \dots, S_B^*) serves as an approximation to the null distribution of S .

The (simulated) bootstrap P -value for a test that rejects the null hypothesis (1) for large values of S is computed as the proportion of (S_1^*, \dots, S_B^*) greater than the observed value of S . Hence, for a given nominal level α ($0 < \alpha < 1$), the bootstrap test rejects \mathcal{H}_0 if the bootstrap P -value does not exceed α . Equivalently, the bootstrap test of level α rejects \mathcal{H}_0 if S exceeds the $(\lfloor (B + 1)(1 - \alpha) \rfloor)$ th largest of (S_1^*, \dots, S_B^*) .

Some remarks about the bootstrap procedure are in order.

(i) The order p of the autoregressive sieve in step S1 may be selected from a suitable range of values by means of the Akaike information criterion (AIC), so as to minimize $\log \hat{\sigma}_p^2 + 2p/n$. Under mild regularity conditions, a data-dependent choice of p based on the AIC is asymptotically efficient (see, inter alia, Shibata (1980), Lee and Karagrigoriou (2001), and Poskitt (2007)), and satisfies the growth conditions on the sieve order that are typically required for the asymptotic validity of the sieve bootstrap for a large class of statistics (Psaradakis (2016)).

(ii) Although least-squares estimates $(\hat{\phi}_{p1}, \dots, \hat{\phi}_{pp}, \hat{\sigma}_p^2)$ of the parameters of the approximating autoregression are used in step S2 to construct X_t^* , asymptotically equivalent estimates, such as those obtained from the empirical Yule–Walker equations, may alternatively be used. The Yule–Walker estimator is theoretically attractive because its use guarantees that the bootstrap

pseudo-observations (X_1^*, \dots, X_n^*) are generated from a causal (bootstrap) autoregressive process, but is known to be significantly biased in small samples compared to the least-squares estimator (see, e.g., Tjøstheim and Paulsen (1983) and Paulsen and Tjøstheim (1985)).

(iii) By requiring ε_t^* in (14) to be normally distributed, the bootstrap pseudo-data $\{X_t^*\}$ are constructed in a way which reflects the normality hypothesis under test even though \mathcal{X} may not satisfy (1). This is important for ensuring that the bootstrap test has reasonable power against departures from \mathcal{H}_0 (see, e.g., Lehmann and Romano (2005, Sec. 15.6)).

(iv) Some variations of the bootstrap procedure may be obtained by varying the way in which the initial values $(X_{-p+1}^*, \dots, X_0^*)$ for the recursion (14) are chosen in step S2. For instance, one possibility is to calculate $(X_{-p+1}^*, \dots, X_0^*)$ from the moving-average representation of the fitted autoregressive model for $X_t - \bar{X}$ (Paparoditis and Streitberg (1992)). Another possibility is to set $X_t^* = X_{t+q}$ for $t \leq 0$, where q is chosen randomly from the set of integers $\{p, p+1, \dots, n\}$ (e.g., Poskitt (2008)). In the sequel, we follow the suggestion of Bühlmann (1997) and set $X_t^* = \bar{X}$ for $t \leq 0$, generate $n + n_0$ bootstrap replicates X_t^* according to (14), with $n_0 = 100$, and then discard the first n_0 replicates to minimize the effect of initial values.

We conclude this section by noting that the linear structure assumed in (11) or (12) may arguably be considered as somewhat restrictive. However, since nonlinear processes with a Gaussian marginal distribution appear to be a rarity (cf. Tong (1990, Sec. 4.2)), the assumption of linear dependence is not perhaps unjustifiable when the objective is to test for marginal normality.

Moreover, the results of Bickel and Bühlmann (1997) indicate that linearity may not be too onerous a requirement, in the sense that the closure (with respect to certain metrics) of the class of causal linear processes is quite large; roughly speaking, for any strictly stationary nonlinear process, there exists another process in the closure of causal linear processes having identical sample paths with probability exceeding 0.36. This also suggests that the autoregressive sieve bootstrap is likely to yield reasonably good approximations within a class of processes larger than that associated with (11) or (12). In fact, Kreiss et al. (2011) have demonstrated that the autoregressive sieve bootstrap is asymptotically valid for a general class of statistics associated with strictly stationary, weakly dependent, regular processes having positive and bounded spectral densities. Such processes can always be represented in the form (11) and (12), with $\{\varepsilon_t\}$ being a strictly stationary sequence of uncorrelated (although not necessarily independent) random variables. Then, the autoregressive coefficients in (11) may also be thought of as the limit, as p tends to infinity, of the coefficients of the best linear predictor (in a mean-square sense) of $X_t - \mu_X$ based on the finite past $(X_{t-1} - \mu_X, \dots, X_{t-p} - \mu_X)$ of length p . The finite-predictor coefficients of \mathcal{X} are uniquely determined for each fixed integer $p \geq 1$ as long as $\sigma_X^2 > 0$ and $\text{Cov}(X_t, X_{t-\tau}) \rightarrow 0$ as $\tau \rightarrow \infty$ (cf. Brockwell and Davis (1991, Sec. 5.1)), and converge to the corresponding infinite-predictor coefficients as $p \rightarrow \infty$ (cf. Pourahmadi (2001, Sec. 7.6), Kreiss et al. (2011)).

4. SIMULATION STUDY

In this section we present and discuss the results of a simulation study examining the finite-sample properties of the normality tests described earlier under various data-generating mechanisms.

4.1 EXPERIMENTAL DESIGN

In the first set of experiments, we examine the performance of normality tests under different patterns of dependence by considering artificial data generated according to the ARMA models

$$\mathbf{M1:} \quad X_t = 0.8X_{t-1} + \varepsilon_t,$$

$$\mathbf{M2:} \quad X_t = 0.6X_{t-1} - 0.5X_{t-2} + \varepsilon_t,$$

$$\mathbf{M3:} \quad X_t = 0.6X_{t-1} + 0.3\varepsilon_{t-1} + \varepsilon_t.$$

Here, and throughout this section, $\{\varepsilon_t\}$ are i.i.d. random variables the common distribution of which is either standard normal (labelled N in the various tables) or generalized lambda with quantile function $Q_\varepsilon(w) = \lambda_1 + (1/\lambda_2)\{w^{\lambda_3} - (1-w)^{\lambda_4}\}$, $0 < w < 1$, standardized to have zero mean and unit variance (see Ramberg and Schmeiser (1974)). The parameter values of the generalized lambda distribution used in the experiments are taken from Bai and Ng (2005) and can be found in Table 1, along with the corresponding coefficients of skewness and kurtosis; the distributions S1–S3 are symmetric, whereas A1–A4 are asymmetric.

In addition, we consider artificial data generated according to the transformation model

$$\mathbf{M4:} \quad X_t = \Phi^{-1}(F_\xi(\xi_t)), \quad \xi_t = \theta|\xi_{t-1}| + \varepsilon_t, \quad \varepsilon_t \sim \mathcal{N}(0, 1), \quad \theta = 0.5,$$

where F_ξ is the distribution function of ξ_t . The process $\{X_t\}$ obtained from the threshold autoregressive process $\{\xi_t\}$ through the composite function $\Phi^{-1} \circ F_\xi$ does not admit the representation (11) or (12) (with respect to i.i.d. innovations), but satisfies the null hypothesis since $X_t \sim \mathcal{N}(0, 1)$ for each t . Note that $\{\xi_t\}$ is strictly stationary with

$$F_\xi(u) = \{2(1 - \theta^2)/\pi\}^{1/2} \int_{-\infty}^u \exp\{-(1 - \theta^2)x^2/2\} \Phi(\theta x) dx, \quad -\infty < u < \infty,$$

for all $|\theta| < 1$ (Anděl and Ranocha (2005)).

The effect of nonlinearity on the properties of the tests is explored further in a second set of experiments by using artificial data from the models

$$\mathbf{M5:} \quad X_t = (0.9X_{t-1} + \varepsilon_t)\mathbb{I}(|X_{t-1}| \leq 1) - (0.3X_{t-1} + 2\varepsilon_t)\mathbb{I}(|X_{t-1}| > 1),$$

$$\mathbf{M6:} \quad X_t = (0.8X_{t-1} + \varepsilon_t)\{1 - \Lambda(X_{t-1})\} - (0.8X_{t-1} + 2\varepsilon_t)\Lambda(X_{t-1}),$$

$$\mathbf{M7:} \quad X_t = \eta_t \varepsilon_t, \quad \eta_t^2 = 0.05 + 0.1X_{t-1}^2 + 0.85\eta_{t-1}^2,$$

$$\mathbf{M8:} \quad X_t = 0.7X_{t-2}\varepsilon_{t-1} + \varepsilon_t,$$

where $\Lambda(x) = 1/(1 + e^{-x})$ is the standard logistic function and $\mathbb{I}(A)$ denotes the indicator of the event A . M5 is a threshold autoregressive model, M6 is a smooth-transition autoregressive model, M7 is a generalized autoregressive conditionally heteroskedastic model, and M8 is a bilinear model. In all four cases, $\{X_t\}$ does not admit the representation (11) or (12); furthermore, the distribution of X_t is non-Gaussian even if ε_t is normally distributed.

For each design point, 1,000 independent realizations of $\{X_t\}$ of length $100 + n$, with $n \in \{100, 200\}$, are generated. The first 100 data points of each realization are then discarded in order to eliminate start-up effects and the remaining n data points are used to compute the value of the test statistics defined in (2)–(10). In the case of bootstrap tests, the order of the autoregressive sieve is determined by minimizing the AIC in the range $1 \leq p \leq \lfloor 10 \log_{10} n \rfloor$, while the number of bootstrap replications is $B = 199$. (We note that using a larger number of bootstrap replications did not change the results substantially. Hall (1986) and Jöckel (1986) provide theoretical explanations of the ability of simulation-based inference procedures to yield good results for relatively small values of the simulation size).

4.2 SIMULATION RESULTS

The Monte Carlo rejection frequencies of normality tests at the 5% significance level ($\alpha = 0.05$) are reported in Tables 2–9. Asymptotic tests (based on JB , BN , LV , BM , and H) rely on critical values from the relevant chi-square distribution; bootstrap tests use critical values obtained by an autoregressive sieve bootstrap procedure. The results over all design points which do not satisfy the null hypothesis are summarized graphically in the form of the box plot of the empirical rejection frequencies shown in Figure 1 (bootstrap tests are indicated by the subscript B).

Inspection of the results in Tables 2–4 (under Gaussian innovations) and in Table 5 reveals that the test based on H suffers from severe level distortion across all four data-generating mechanisms when asymptotic critical values are used. Among the remaining asymptotic tests, LV has an overall advantage under the null hypothesis for both of the sample sizes considered. The BN and BM tests tend to be too liberal and, rather surprisingly, do not perform substantially better than the JB test, which relies on the assumption of i.i.d. observations. A possible explanation for the unsatisfactory level performance of the tests based on BN and BM may lie with the kernel estimators of the relevant long-run covariance matrices that are used in their construction. Inference procedures based on such estimators are widely reported to have poor small-sample properties, and related tests are often found to exhibit substantial level distortions in a variety of settings (see, e.g., den Haan and Levin (1997), Müller (2014)). As expected perhaps, bootstrap tests are generally more successful than asymptotic tests at



controlling the discrepancy between the exact and nominal probabilities of Type I error. The empirical rejection frequencies of bootstrap tests are insignificantly different from the nominal 0.05 value in the vast majority of cases.

The results in Tables 2–4 (under non-Gaussian innovations) and in Tables 6–9 show that the bootstrap versions of the JB and LV tests tend to outperform all other tests in terms of empirical power, albeit only marginally in some cases, regardless of the dependence structure in the data and the distribution of the innovations. In particular, as can be easily seen in Figure 1, for processes with a non-Gaussian marginal distribution, the bootstrap JB and LV tests have the highest average rejection frequencies (indicated by black diamonds) across all tests, and smaller interquartile range (edges of coloured areas) than the asymptotic LV test. However, keeping in mind computational aspects and level accuracy, the latter test offers an attractive alternative to bootstrap tests. Among tests based on the empirical distribution function, which are also competitive in terms of power, the AD and CM tests tend to have a slight advantage over the KS test, and perform quite similarly to the EP test based on the empirical characteristic function. The rejection frequencies of the asymptotic and bootstrap BN and BM tests have distributions which are highly positively skewed (cf. Figure 1), which means that the tests are powerful only for some design points. Rather unsurprisingly, the rejection frequencies of tests improve with increasing skewness and leptokurtosis in the innovation distribution, as well as with an increasing sample size. It is worth noting that, although the asymptotic versions of some tests may appear in some cases to have similar or even higher empirical power than the corresponding bootstrap tests, such comparisons are not straightforward because asymptotic tests do not generally control the probability of Type I error as well as bootstrap tests do. (The asymptotic test based on H is not included in Figure 1 because of its excessive level distortion).

Finally, the simulation results reveal that deviations from the linearity assumptions which underlie the autoregressive sieve bootstrap procedure do not have an adverse effect on the properties of bootstrap tests. Such tests generally work well even for data that are generated by processes which are not representable as in (11) or (12). As can be seen in Table 5, in the case of artificial time series from M4, the marginal distribution of which is Gaussian, most bootstrap tests have rejection frequencies that do not differ substantially from the nominal level (the AD and CM tests have a tendency to over-reject). Similarly, as can be seen in Tables 6–9, the bootstrap versions of tests other than BN and BM have high rejection frequencies for data with a non-Gaussian marginal distribution generated according to the non-linear models M5–M8.

5. SUMMARY

This paper has considered the problem of testing for normality of the one-dimensional marginal distribution of a strictly stationary and weakly dependent stochastic process. We have ex-



amined the properties of nine normality tests, only some of which have been designed to be robust with respect to dependence in the data. Since conventional large-sample approximations to the null distributions of some of the test statistics are either unknown or inaccurate under dependence, we have explored how an autoregressive sieve bootstrap procedure may be used to obtain P -values and/or critical values for the tests. An extensive Monte Carlo study has revealed that the bootstrap version of the classical skewness–kurtosis test provides the best overall performance across the asymptotic and bootstrap tests investigated. The Lobato–Velasco modification of the cumulant-based test is a good alternative among tests that rely on asymptotic critical values.



REFERENCES

- Andrews, D. W. K. (1991). Heteroskedasticity and autocorrelation consistent covariance matrix estimation. *Econometrica* 59, 817–858.
- Anděl, J. and P. Ranocha (2005). Stationary distribution of absolute autoregression. *Kybernetika* 41, 735–742.
- Babu, G. J. and C. R. Rao (2004). Goodness-of-fit tests when parameters are estimated. *Sankhyā* 66, 63–74.
- Bai, J. and S. Ng (2005). Tests for skewness, kurtosis, and normality for time series data. *Journal of Business and Economic Statistics* 23, 49–60.
- Baringhaus, L., R. Danchke, and N. Henze (1989). Recent and classical tests for normality - a comparative study. *Communications in Statistics – Simulation and Computation* 18, 363–379.
- Baringhaus, L. and N. Henze (1988). A consistent test for multivariate normality based on the empirical characteristic function. *Metrika* 35, 339–348.
- Berg, A., E. Paparoditis, and D. N. Politis (2010). A bootstrap test for time series linearity. *Journal of Statistical Planning and Inference* 140, 3841–3857.
- Bickel, P. J. and P. Bühlmann (1997). Closure of linear processes. *Journal of Theoretical Probability* 10, 445–479.
- Bontemps, C. and N. Meddahi (2005). Testing normality: a GMM approach. *Journal of Econometrics* 124, 149–186.
- Bowman, K. O. and L. R. Shenton (1975). Omnibus test contours for departures from normality based on $\sqrt{b_1}$ and b_2 . *Biometrika* 62, 243–250.
- Brockwell, P. J. and R. A. Davis (1991). *Time Series: Theory and Methods*. New York: 2nd Edition, Springer.
- Bühlmann, P. (1997). Sieve bootstrap for time series. *Bernoulli* 3, 123–148.
- Choi, E. and P. Hall (2000). Bootstrap confidence regions constructed from autoregressions of arbitrary order. *Journal of the Royal Statistical Society, Ser. B* 62, 461–477.
- Cotter, J. (2007). Varying the VaR for unconditional and conditional environments. *Journal of International Money and Finance* 26, 1338–1354.
- den Haan, W. J. and A. Levin (1997). A practitioner's guide to robust covariance matrix estimation. In G. S. Maddala and C. R. Rao (Eds.), *Handbook of Statistics: Robust Inference*, Volume 15, pp. 299–342. Amsterdam: North-Holland.



- Durbin, J. (1973). Weak convergence of the sample distribution function when parameters are estimated. *Annals of Statistics* 1, 279–290.
- Epps, T. W. (1987). Testing that a stationary time series is Gaussian. *Annals of Statistics* 15, 1683–1698.
- Epps, T. W. and L. B. Pulley (1983). A test for normality based on the empirical characteristic function. *Biometrika* 70, 723–726.
- Hall, P. (1986). On the number of bootstrap simulations required to construct a confidence interval. *Annals of Statistics* 14, 1453–1462.
- Hinich, M. J. (1982). Testing for Gaussianity and linearity of a stationary time series. *Journal of Time Series Analysis* 3, 169–176.
- Jarque, C. M. and A. K. Bera (1987). A test for normality of observations and regression residuals. *International Statistical Review* 55, 163–172.
- Jiménez-Gamero, M. D., J. Muñoz García, and R. Pino-Mejías (2003). Bootstrapping parameter estimated degenerate U and V statistics. *Statistics and Probability Letters* 61, 61–70.
- Jöckel, K.-H. (1986). Finite-sample properties and asymptotic efficiency of Monte Carlo tests. *Annals of Statistics* 14, 336–347.
- Kilian, L. and U. Demiroglu (2000). Residual-based tests for normality in autoregressions: asymptotic theory and simulation evidence. *Journal of Business and Economic Statistics* 18, 40–50.
- Kojadinovic, I. and J. Yan (2012). Goodness-of-fit testing based on a weighted bootstrap: a fast large-sample alternative to the parametric bootstrap. *Canadian Journal of Statistics* 40, 480–500.
- Kreiss, J.-P. (1992). Bootstrap procedures for $AR(\infty)$ processes. In K.-H. Jöckel, G. Rothe, and W. Sendler (Eds.), *Bootstrapping and Related Techniques*, pp. 107–113. Heidelberg: Springer-Verlag.
- Kreiss, J.-P., E. Paparoditis, and D. N. Politis (2011). On the range of validity of the autoregressive sieve bootstrap. *Annals of Statistics* 39, 2103–2130.
- Lee, S. and A. Karagrigoriou (2001). An asymptotically optimal selection of the order of a linear process. *Sankhyā, Ser. A* 63, 93–106.
- Lehmann, E. L. and J. P. Romano (2005). *Testing Statistical Hypotheses*. New York: 3rd Edition, Springer.
- Leucht, A. (2012). Degenerate U - and V -statistics under weak dependence: asymptotic theory and bootstrap consistency. *Bernoulli* 18, 552–585.



- Leucht, A. and M. H. Neumann (2009). Consistency of general bootstrap methods for degenerate U -type and V -type statistics. *Journal of Multivariate Analysis* 100, 1622–1633.
- Leucht, A. and M. H. Neumann (2013). Degenerate U - and V -statistics under ergodicity: asymptotics, bootstrap and applications in statistics. *Annals of the Institute of Statistical Mathematics* 65, 349–386.
- Lobato, I. N. and C. Velasco (2004). A simple test of normality for time series. *Econometric Theory* 20, 671–689.
- Moulines, E. and K. Choukri (1996). Time-domain procedures for testing that a stationary time-series is Gaussian. *IEEE Transactions on Signal Processing* 44, 2010–2025.
- Müller, U. K. (2014). HAC corrections for strongly autocorrelated time series. *Journal of Business and Economic Statistics* 32, 311–322.
- Nusrat, J. and J. L. Harvill (2008). Bispectral-based goodness-of-fit tests of Gaussianity and linearity of stationary time series. *Communications in Statistics – Theory and Methods* 37, 3216–3227.
- Paparoditis, E. (1996). Bootstrapping autoregressive and moving average parameter estimates of infinite order vector autoregressive processes. *Journal of Multivariate Analysis* 57, 277–296.
- Paparoditis, E. and B. Streitberg (1992). Order identification statistics in stationary autoregressive moving-average models: vector autocorrelations and the bootstrap. *Journal of Time Series Analysis* 13, 415–434.
- Paulsen, J. and D. Tjøstheim (1985). On the estimation of residual variance and order in autoregressive time series. *Journal of the Royal Statistical Society, Ser. B* 47, 216–228.
- Poskitt, D. S. (2007). Autoregressive approximation in nonstandard situations: the fractionally integrated and non-invertible cases. *Annals of the Institute of Statistical Mathematics* 59, 697–725.
- Poskitt, D. S. (2008). Properties of the sieve bootstrap for fractionally integrated and non-invertible processes. *Journal of Time Series Analysis* 29, 224–250.
- Pourahmadi, M. (2001). *Foundations of Time Series Analysis and Prediction Theory*. New York: Wiley.
- Psaradakis, Z. (2016). Using the bootstrap to test for symmetry under unknown dependence. *Journal of Business and Economic Statistics* 34, 406–415.
- Psaradakis, Z. and M. Vávra (2017). A distance test of normality for a wide class of stationary processes. *Econometrics and Statistics* 2, 50–60.




- Ramberg, J. S. and B. W. Schmeiser (1974). An approximate method for generating asymmetric random variables. *Communications of the ACM* 17, 78–82.
- Romão, X., R. Delgado, and A. Costa (2010). An empirical power comparison of univariate goodness-of-fit tests for normality. *Journal of Statistical Computation and Simulation* 80, 545–591.
- Shibata, R. (1980). Asymptotically efficient selection of the order of the model for estimating parameters of a linear process. *Annals of Statistics* 8, 147–164.
- Stein, C. (1972). A bound for the error in the normal approximation to the distribution of a sum of dependent random variables. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Volume 2: Probability Theory*, pp. 583–602. Berkeley: University of California Press.
- Stephens, M. A. (1976). Asymptotic results for goodness-of-fit statistics with unknown parameters. *Annals of Statistics* 4, 357–369.
- Stute, W., W. Gonzáles Manteiga, and M. Presedo Quindimil (1993). Bootstrap based goodness-of-fit-tests. *Metrika* 40, 243–256.
- Thode, H. C. (2002). *Testing for Normality*. New York: Marcel Dekker.
- Tjøstheim, D. and J. Paulsen (1983). Bias of some commonly-used time series estimates. *Biometrika* 70, 389–399; Corrigendum (1984), 71, 656.
- Tong, H. (1990). *Non-linear Time Series: A Dynamical System Approach*. Oxford: Oxford University Press.
- Yap, B. W. and C. H. Sim (2011). Comparisons of various types of normality tests. *Journal of Statistical Computation and Simulation* 81, 2141–2155.

Table 1: Innovation Distributions

	λ_1	λ_2	λ_3	λ_4	skewness	kurtosis
N	–	–	–	–	0.0	3.0
S1	0.000000	-1.000000	-0.080000	-0.080000	0.0	6.0
S2	0.000000	-0.397912	-0.160000	-0.160000	0.0	11.6
S3	0.000000	-1.000000	-0.240000	-0.240000	0.0	126.0
A1	0.000000	-1.000000	-0.007500	-0.030000	1.5	7.5
A2	0.000000	-1.000000	-0.100900	-0.180200	2.0	21.1
A3	0.000000	-1.000000	-0.001000	-0.130000	3.2	23.8
A4	0.000000	-1.000000	-0.000100	-0.170000	3.8	40.7

Figure 1: Empirical Rejection Frequencies of Normality Tests: Power



Note: The top and bottom of each blue box indicates the 25th and 75th percentile, respectively, of the empirical rejection frequencies, the black diamond indicates the mean value, and the whiskers indicate the 10th and 90th percentiles.

Table 2: Empirical Rejection Frequencies of Normality Tests Under M1

sample	distr.	Asymptotic Tests					Bootstrap Tests								
		<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>AD</i>	<i>CM</i>	<i>KS</i>	<i>EP</i>
<i>n</i> = 100	N	0.09	0.08	0.02	0.07	0.31	0.04	0.05	0.04	0.05	0.03	0.04	0.04	0.04	0.05
	S1	0.22	0.05	0.10	0.04	0.48	0.16	0.03	0.15	0.04	0.13	0.11	0.11	0.11	0.11
	S2	0.32	0.09	0.17	0.09	0.60	0.25	0.06	0.23	0.07	0.24	0.21	0.21	0.18	0.19
	S3	0.44	0.06	0.29	0.06	0.69	0.38	0.04	0.35	0.04	0.34	0.29	0.27	0.23	0.28
	A1	0.39	0.13	0.22	0.11	0.68	0.30	0.09	0.29	0.09	0.31	0.24	0.22	0.18	0.26
	A2	0.46	0.12	0.29	0.12	0.72	0.37	0.08	0.35	0.09	0.37	0.33	0.33	0.28	0.34
	A3	0.74	0.34	0.49	0.34	0.93	0.64	0.23	0.60	0.27	0.64	0.57	0.56	0.47	0.61
	A4	0.81	0.34	0.59	0.34	0.94	0.73	0.24	0.70	0.28	0.72	0.68	0.64	0.55	0.71
<i>n</i> = 200	N	0.16	0.08	0.04	0.08	0.31	0.07	0.06	0.06	0.06	0.04	0.06	0.05	0.05	0.07
	S1	0.34	0.05	0.16	0.03	0.53	0.20	0.03	0.20	0.02	0.19	0.11	0.10	0.07	0.11
	S2	0.51	0.06	0.29	0.07	0.66	0.35	0.04	0.35	0.04	0.30	0.25	0.24	0.21	0.25
	S3	0.63	0.07	0.45	0.08	0.74	0.52	0.05	0.51	0.06	0.44	0.39	0.38	0.34	0.38
	A1	0.68	0.32	0.41	0.34	0.82	0.51	0.25	0.48	0.27	0.40	0.45	0.42	0.35	0.46
	A2	0.67	0.21	0.45	0.21	0.80	0.52	0.14	0.52	0.16	0.46	0.45	0.44	0.38	0.46
	A3	0.96	0.67	0.81	0.67	0.98	0.86	0.53	0.85	0.55	0.84	0.86	0.81	0.74	0.88
	A4	0.99	0.68	0.87	0.68	1.00	0.93	0.56	0.92	0.60	0.91	0.92	0.88	0.82	0.93



Table 3: Empirical Rejection Frequencies of Normality Tests Under M2

sample	distr.	Asymptotic Tests					Bootstrap Tests								
		<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>AD</i>	<i>CM</i>	<i>KS</i>	<i>EP</i>
<i>n</i> = 100	N	0.03	0.07	0.03	0.06	0.28	0.05	0.04	0.05	0.05	0.04	0.04	0.03	0.04	0.03
	S1	0.33	0.06	0.32	0.05	0.46	0.38	0.04	0.36	0.04	0.15	0.24	0.19	0.15	0.24
	S2	0.47	0.07	0.45	0.06	0.55	0.50	0.03	0.49	0.04	0.23	0.38	0.34	0.28	0.39
	S3	0.68	0.07	0.67	0.07	0.66	0.71	0.03	0.70	0.03	0.36	0.55	0.52	0.41	0.55
	A1	0.64	0.39	0.66	0.37	0.68	0.69	0.28	0.70	0.31	0.29	0.63	0.57	0.52	0.69
	A2	0.66	0.14	0.65	0.15	0.68	0.70	0.09	0.70	0.10	0.35	0.55	0.52	0.42	0.59
	A3	0.97	0.62	0.97	0.62	0.92	0.98	0.48	0.98	0.53	0.68	0.98	0.97	0.91	0.99
	A4	0.97	0.56	0.97	0.55	0.95	0.99	0.43	0.99	0.47	0.76	0.98	0.96	0.91	0.99
<i>n</i> = 200	N	0.05	0.11	0.05	0.09	0.27	0.05	0.06	0.05	0.06	0.04	0.05	0.06	0.06	0.06
	S1	0.53	0.04	0.51	0.05	0.51	0.54	0.02	0.53	0.02	0.16	0.32	0.26	0.19	0.32
	S2	0.74	0.08	0.73	0.08	0.67	0.76	0.03	0.73	0.03	0.29	0.55	0.50	0.41	0.59
	S3	0.87	0.12	0.86	0.11	0.72	0.88	0.03	0.87	0.03	0.46	0.75	0.71	0.61	0.77
	A1	0.97	0.77	0.96	0.76	0.76	0.96	0.61	0.96	0.64	0.38	0.92	0.90	0.78	0.95
	A2	0.91	0.28	0.91	0.29	0.78	0.91	0.17	0.91	0.18	0.48	0.84	0.80	0.69	0.86
	A3	1.00	0.87	1.00	0.87	0.99	1.00	0.78	1.00	0.80	0.88	1.00	1.00	1.00	1.00
	A4	1.00	0.83	1.00	0.83	0.99	1.00	0.74	1.00	0.77	0.90	1.00	1.00	1.00	1.00



Table 4: Empirical Rejection Frequencies of Normality Tests Under M3

sample	distr.	Asymptotic Tests					Bootstrap Tests								
		<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>AD</i>	<i>CM</i>	<i>KS</i>	<i>EP</i>
<i>n</i> = 100	N	0.08	0.08	0.03	0.07	0.33	0.05	0.04	0.04	0.05	0.05	0.04	0.03	0.03	0.05
	S1	0.27	0.04	0.18	0.04	0.52	0.22	0.02	0.21	0.03	0.14	0.16	0.13	0.11	0.15
	S2	0.41	0.08	0.32	0.07	0.57	0.38	0.05	0.37	0.05	0.21	0.28	0.26	0.22	0.27
	S3	0.51	0.08	0.43	0.07	0.65	0.47	0.05	0.47	0.05	0.34	0.39	0.37	0.32	0.38
	A1	0.59	0.28	0.42	0.28	0.74	0.52	0.22	0.49	0.23	0.29	0.47	0.43	0.36	0.50
	A2	0.53	0.16	0.43	0.13	0.71	0.49	0.10	0.48	0.10	0.36	0.43	0.40	0.37	0.41
	A3	0.91	0.58	0.82	0.58	0.92	0.87	0.45	0.88	0.49	0.68	0.89	0.86	0.77	0.91
	A4	0.96	0.56	0.88	0.54	0.96	0.92	0.43	0.92	0.47	0.76	0.94	0.90	0.85	0.94
<i>n</i> = 200	N	0.10	0.10	0.04	0.09	0.34	0.05	0.06	0.05	0.05	0.05	0.05	0.05	0.06	0.06
	S1	0.45	0.06	0.30	0.06	0.52	0.33	0.04	0.34	0.04	0.17	0.19	0.18	0.15	0.19
	S2	0.64	0.08	0.52	0.08	0.64	0.53	0.04	0.54	0.04	0.29	0.37	0.36	0.28	0.37
	S3	0.78	0.12	0.69	0.11	0.77	0.72	0.07	0.72	0.07	0.46	0.58	0.55	0.47	0.59
	A1	0.87	0.59	0.75	0.60	0.81	0.78	0.44	0.76	0.46	0.45	0.76	0.72	0.61	0.77
	A2	0.83	0.29	0.70	0.28	0.81	0.75	0.18	0.74	0.20	0.51	0.69	0.66	0.59	0.71
	A3	1.00	0.94	0.99	0.94	0.99	1.00	0.86	1.00	0.88	0.87	1.00	1.00	0.98	1.00
	A4	1.00	0.92	1.00	0.92	1.00	1.00	0.85	1.00	0.87	0.93	1.00	1.00	0.99	1.00



Table 5: Empirical Rejection Frequencies of Normality Tests Under M4

sample	distr.	Asymptotic Tests					Bootstrap Tests								
		<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>AD</i>	<i>CM</i>	<i>KS</i>	<i>EP</i>
<i>n</i> = 100	N	0.02	0.13	0.02	0.11	0.23	0.04	0.07	0.03	0.07	0.06	0.09	0.08	0.03	0.08
<i>n</i> = 200	N	0.02	0.12	0.02	0.11	0.25	0.04	0.06	0.03	0.06	0.06	0.11	0.10	0.03	0.07



Table 6: Empirical Rejection Frequencies of Normality Tests Under M5

sample	distr.	Asymptotic Tests					Bootstrap Tests								
		<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>AD</i>	<i>CM</i>	<i>KS</i>	<i>EP</i>
<i>n</i> = 100	N	0.35	0.08	0.34	0.07	0.40	0.37	0.03	0.37	0.04	0.13	0.26	0.25	0.19	0.25
	S1	0.79	0.12	0.78	0.13	0.59	0.79	0.07	0.80	0.08	0.33	0.69	0.65	0.54	0.70
	S2	0.86	0.14	0.86	0.14	0.65	0.87	0.08	0.87	0.11	0.43	0.79	0.76	0.62	0.78
	S3	0.92	0.15	0.90	0.14	0.71	0.92	0.09	0.92	0.10	0.49	0.88	0.86	0.77	0.89
	A1	0.78	0.07	0.78	0.07	0.65	0.80	0.03	0.79	0.04	0.39	0.66	0.59	0.52	0.64
	A2	0.81	0.11	0.80	0.12	0.64	0.83	0.07	0.83	0.08	0.42	0.75	0.70	0.60	0.75
	A3	0.92	0.07	0.91	0.07	0.79	0.92	0.04	0.92	0.05	0.58	0.84	0.76	0.69	0.80
	A4	0.91	0.08	0.90	0.08	0.78	0.91	0.05	0.91	0.06	0.60	0.86	0.77	0.70	0.83
<i>n</i> = 200	N	0.52	0.13	0.51	0.13	0.42	0.52	0.05	0.52	0.06	0.16	0.42	0.40	0.29	0.43
	S1	0.94	0.25	0.93	0.27	0.66	0.93	0.12	0.94	0.13	0.38	0.90	0.87	0.76	0.91
	S2	0.98	0.24	0.98	0.24	0.74	0.98	0.12	0.98	0.13	0.48	0.96	0.96	0.89	0.97
	S3	0.99	0.25	0.99	0.25	0.82	0.99	0.12	0.99	0.13	0.60	0.99	0.98	0.96	0.99
	A1	0.95	0.16	0.95	0.16	0.73	0.95	0.06	0.95	0.08	0.49	0.90	0.83	0.75	0.88
	A2	0.98	0.19	0.98	0.19	0.79	0.98	0.08	0.98	0.09	0.55	0.97	0.96	0.90	0.97
	A3	0.99	0.15	0.99	0.16	0.88	0.99	0.07	0.99	0.07	0.72	0.97	0.93	0.91	0.97
	A4	0.99	0.13	0.99	0.13	0.90	0.99	0.04	0.99	0.06	0.75	0.99	0.97	0.96	0.98



Table 7: Empirical Rejection Frequencies of Normality Tests Under M6

sample	distr.	Asymptotic Tests					Bootstrap Tests								
		<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>AD</i>	<i>CM</i>	<i>KS</i>	<i>EP</i>
<i>n</i> = 100	N	0.86	0.24	0.85	0.24	0.86	0.87	0.19	0.87	0.19	0.68	0.91	0.92	0.86	0.88
	S1	0.95	0.25	0.95	0.24	0.88	0.95	0.16	0.95	0.17	0.71	0.97	0.96	0.93	0.96
	S2	0.97	0.24	0.97	0.23	0.88	0.97	0.15	0.97	0.17	0.76	0.98	0.99	0.96	0.98
	S3	0.98	0.25	0.98	0.25	0.92	0.98	0.17	0.98	0.18	0.79	0.99	0.99	0.97	0.99
	A1	0.93	0.33	0.93	0.32	0.87	0.93	0.20	0.93	0.21	0.69	0.92	0.92	0.89	0.92
	A2	0.95	0.27	0.95	0.26	0.89	0.95	0.16	0.95	0.18	0.73	0.96	0.95	0.91	0.97
	A3	1.00	0.72	1.00	0.72	0.94	1.00	0.62	1.00	0.66	0.81	1.00	0.99	0.97	1.00
	A4	1.00	0.72	1.00	0.71	0.97	1.00	0.60	1.00	0.64	0.89	1.00	1.00	0.99	1.00
<i>n</i> = 200	N	0.99	0.31	0.99	0.31	0.96	0.99	0.17	0.99	0.19	0.81	1.00	1.00	0.99	0.99
	S1	1.00	0.32	1.00	0.31	0.95	1.00	0.17	1.00	0.18	0.87	1.00	1.00	1.00	1.00
	S2	1.00	0.31	1.00	0.31	0.98	1.00	0.17	1.00	0.18	0.88	1.00	1.00	1.00	1.00
	S3	1.00	0.21	1.00	0.20	0.98	1.00	0.09	1.00	0.10	0.93	1.00	1.00	1.00	1.00
	A1	1.00	0.71	1.00	0.70	0.96	1.00	0.53	1.00	0.55	0.84	0.99	0.99	0.99	1.00
	A2	1.00	0.38	1.00	0.37	0.98	1.00	0.22	1.00	0.25	0.89	1.00	1.00	1.00	1.00
	A3	1.00	0.88	1.00	0.89	1.00	1.00	0.81	1.00	0.82	0.96	1.00	1.00	1.00	1.00
	A4	1.00	0.89	1.00	0.88	1.00	1.00	0.82	1.00	0.84	0.97	1.00	1.00	1.00	1.00



Table 8: Empirical Rejection Frequencies of Normality Tests Under M7

sample	distr.	Asymptotic Tests					Bootstrap Tests								
		<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>AD</i>	<i>CM</i>	<i>KS</i>	<i>EP</i>
<i>n</i> = 100	N	0.09	0.09	0.09	0.10	0.39	0.07	0.05	0.11	0.05	0.21	0.07	0.07	0.06	0.08
	S1	0.64	0.11	0.64	0.11	0.58	0.67	0.05	0.66	0.06	0.36	0.52	0.50	0.37	0.53
	S2	0.79	0.14	0.79	0.14	0.63	0.80	0.07	0.80	0.09	0.44	0.73	0.70	0.61	0.75
	S3	0.92	0.17	0.91	0.16	0.71	0.92	0.09	0.92	0.11	0.56	0.90	0.88	0.81	0.90
	A1	0.94	0.76	0.94	0.75	0.70	0.95	0.63	0.95	0.64	0.51	0.97	0.96	0.91	0.98
	A2	0.92	0.29	0.92	0.29	0.74	0.92	0.15	0.92	0.18	0.55	0.90	0.88	0.83	0.92
	A3	1.00	0.91	1.00	0.90	0.93	1.00	0.83	1.00	0.85	0.82	1.00	1.00	1.00	1.00
	A4	1.00	0.92	1.00	0.90	0.95	1.00	0.84	1.00	0.86	0.86	1.00	1.00	1.00	1.00
<i>n</i> = 200	N	0.16	0.07	0.16	0.06	0.62	0.17	0.02	0.17	0.02	0.36	0.11	0.08	0.07	0.10
	S1	0.90	0.22	0.90	0.22	0.78	0.90	0.08	0.89	0.10	0.59	0.82	0.79	0.66	0.84
	S2	0.97	0.27	0.97	0.27	0.84	0.98	0.12	0.98	0.14	0.70	0.96	0.95	0.92	0.96
	S3	1.00	0.28	1.00	0.27	0.93	1.00	0.11	1.00	0.13	0.76	0.99	0.99	0.98	0.99
	A1	1.00	0.91	1.00	0.91	0.89	1.00	0.83	1.00	0.83	0.74	1.00	1.00	0.99	1.00
	A2	1.00	0.45	1.00	0.45	0.90	0.99	0.28	0.99	0.30	0.78	0.99	0.99	0.97	1.00
	A3	1.00	0.96	1.00	0.97	0.98	1.00	0.93	1.00	0.94	0.93	1.00	1.00	1.00	1.00
	A4	1.00	0.96	1.00	0.96	0.99	1.00	0.93	1.00	0.93	0.95	1.00	1.00	1.00	1.00



Table 9: Empirical Rejection Frequencies of Normality Tests Under M8

sample	distr.	Asymptotic Tests					Bootstrap Tests								
		<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>JB</i>	<i>BN</i>	<i>LV</i>	<i>BM</i>	<i>H</i>	<i>AD</i>	<i>CM</i>	<i>KS</i>	<i>EP</i>
<i>n</i> = 100	N	0.34	0.06	0.34	0.04	0.61	0.36	0.02	0.36	0.02	0.37	0.23	0.21	0.15	0.24
	S1	0.63	0.08	0.63	0.08	0.69	0.66	0.05	0.66	0.06	0.48	0.56	0.54	0.45	0.58
	S2	0.75	0.11	0.75	0.11	0.75	0.77	0.08	0.76	0.08	0.53	0.72	0.69	0.58	0.73
	S3	0.82	0.14	0.82	0.13	0.75	0.84	0.08	0.84	0.10	0.57	0.80	0.77	0.71	0.80
	A1	0.79	0.23	0.79	0.23	0.78	0.81	0.16	0.81	0.17	0.61	0.79	0.76	0.68	0.80
	A2	0.80	0.17	0.80	0.18	0.77	0.82	0.09	0.82	0.11	0.59	0.80	0.78	0.67	0.80
	A3	0.98	0.45	0.98	0.44	0.89	0.97	0.36	0.97	0.36	0.75	0.99	0.99	0.97	0.99
	A4	0.99	0.43	0.99	0.42	0.93	0.99	0.33	0.99	0.34	0.82	0.99	0.99	0.98	1.00
<i>n</i> = 200	N	0.58	0.05	0.57	0.06	0.76	0.58	0.02	0.58	0.02	0.50	0.34	0.30	0.22	0.37
	S1	0.88	0.17	0.88	0.16	0.85	0.89	0.07	0.89	0.08	0.65	0.83	0.79	0.69	0.83
	S2	0.96	0.22	0.96	0.22	0.87	0.96	0.09	0.96	0.10	0.70	0.95	0.93	0.88	0.95
	S3	0.99	0.22	0.99	0.22	0.91	0.99	0.09	0.99	0.11	0.78	0.99	0.98	0.94	0.99
	A1	0.98	0.41	0.98	0.41	0.87	0.97	0.24	0.97	0.25	0.71	0.98	0.98	0.92	0.98
	A2	0.98	0.28	0.98	0.29	0.90	0.98	0.14	0.98	0.16	0.77	0.98	0.97	0.95	0.98
	A3	1.00	0.58	1.00	0.56	0.98	1.00	0.43	1.00	0.45	0.91	1.00	1.00	1.00	1.00
	A4	1.00	0.58	1.00	0.57	0.98	1.00	0.42	1.00	0.43	0.93	1.00	1.00	1.00	1.00

